



TEBUCONAZOLE | GROUP 3 | FUNGICIDE

Axill Solutions Teb 3.6SC

For control of specified diseases on various crops; golf course turf; field, nursery and container ornamentals; and ornamentals in commercial and residential landscapes.

ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole

-1-ethanol.....	40.1%
OTHER INGREDIENTS	59.9%
TOTAL	100.0%

Contains 3.6 pounds tebuconazole per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of an emergency involving this product, call CHEMTREC at 1-800-424-9300 .	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	

See label booklet for First Aid, additional Precautionary Statements, and Directions for Use including Storage and Disposal Instructions.

EPA Reg. No. 93809-13

EPA Est. No. 84154-CHN-001

Manufactured for:

Axill Solutions, LLC
422 Jasmine Way
Roseburg, OR 97471

NET CONTENTS: 2.5 Gallons

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Protective eyewear (goggles, safety glasses or face shield)
- Chemical-resistant gloves, made of any waterproof material (barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, and viton)
- Shoes plus socks

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: Axill Solutions Teb 3.6SC may contaminate water through drift of spray in wind. Axill Solutions Teb 3.6SC has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of Axill Solutions Teb 3.6SC will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with oxidizing agents. Hazardous chemical reactions may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls
- Protective eyewear
- Chemical-resistant gloves made of any waterproof material (barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, and viton)
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Golf Course Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

PRODUCT USE INFORMATION

Read the entire CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY before using this product.

[Aerial application Prohibited in New York State.]

Spray Volume: Axill Solutions Teb 3.6SC may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Chemigation: Apply Axill Solutions Teb 3.6SC through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply Axill Solutions Teb 3.6SC only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. **DO NOT** apply Axill Solutions Teb 3.6SC through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration contact State Extension Service specialists, equipment manufacturers or other experts. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments as needed.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and

connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, for example a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add labeled amount of Axill Solutions Teb 3.6SC into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Axill Solutions Teb 3.6SC must be thoroughly dispersed prior to the addition of other materials. **DO NOT** tank mix with other products containing a prohibition against tank mixing.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Compatibility: To determine the compatibility of Axill Solutions Teb 3.6SC with other products, the following procedure must be followed: Pour the specified proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must select nozzles and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- If the windspeed is 10 miles per hour or less, applicators must use ½ swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use ¾ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor blade diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications

- **DO NOT** release spray at a height greater than 4 feet above the ground or crop canopy.
- For golf course applications, **DO NOT** release spray at a height greater than 2 feet above the ground.
- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications:

- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- Adjust Nozzles - Follow nozzle manufacturers directions for setting up nozzles. To reduce fine droplets, orient nozzles parallel with the airflow in flight.
- BOOM HEIGHT - Ground Boom
For ground equipment, the boom should remain level with the crop and have minimal bounce.
- RELEASE HEIGHT - Aircraft
Higher release heights increase the potential for spray drift.
- SHIELDED SPRAYERS
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.
- WIND
Drift potential generally increases with wind speed. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

Resistance Management

For resistance management, Axill Solutions Teb 3.6SC contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to Axill Solutions Teb 3.6SC and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Follow appropriate resistance-management strategies.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Axill Solutions Teb 3.6SC or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

- When possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM guidance for specific crops and pathogens.
- For further information or to report suspected resistance contact Axill Solutions, LLC. You can also contact your pesticide distributor or university extension specialist to report resistance.

ROTATIONAL CROPS RESTRICTIONS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

DISEASE CONTROL IN CROPS

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Asparagus	Rust (<i>Puccinia</i> spp.)	4 to 6 fl. oz. per acre (0.11 to 0.17 lb ai per acre)	<p>Apply Axill Solutions Teb 3.6SC as a foliar spray to the developing ferns after harvest of spears is completed. Applications may be made using ground or aerial application equipment. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl. oz. of Axill Solutions Teb 3.6SC per acre (0.11 lb ai – 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher specified rate. Repeat applications on a 14-day interval as necessary to maintain control of rust.</p> <p>For optimum disease control, tank mix the lowest specified rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Axill Solutions Teb 3.6SC with other DMI fungicides may lead to resistance.</p>
<p>Restrictions:</p> <p>DO NOT apply to harvestable spears.</p> <p>DO NOT make more than three foliar applications per year (18 fl. oz./acre or 0.51 lb. ai/acre). A 50 foot spray drift buffer zone is required for all aerial applications.</p> <p>Maximum Single Application Rate: 6 fl. oz. (0.17 lb ai) per acre Axill Solutions Teb 3.6SC Maximum Annual Application Rate: 18 fl. oz. (0.51 lb ai) per acre per year Axill Solutions Teb 3.6SC Minimum Retreatment Interval (RTI): 14 days</p>			
<p>Pre-Harvest Interval (PHI): DO NOT apply within 100 days of harvest in California and within 180 days of harvest in all other states. Restricted-entry interval (REI) = 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Barley	Rust (<i>Puccinia</i> spp.) Head blight (<i>Fusarium</i> spp.) – Suppression	4 fl. oz. per acre (0.11 lb ai per acre)	<p>Apply Axill Solutions Teb 3.6SC in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. Observe barley fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p>Application timing directions: Rusts: Apply Axill Solutions Teb 3.6SC at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Axill Solutions Teb 3.6SC for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants. For optimum disease control, tank mix the lowest specified rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p>
<p>Restrictions:</p> <p>Straw cut after harvest may be fed or used for bedding.</p> <p>Grazing livestock or feeding of green forage is permitted 6 or more days after application of Axill Solutions Teb 3.6SC.</p> <p>Maximum Single Application Rate: 4 fl. oz. (0.11 lb ai) per acre Axill Solutions Teb 3.6SC. DO NOT make more than one application per year.</p> <p>Pre-Harvest Interval (PHI): DO NOT apply within 30 days of harvest.</p> <p>Restricted-entry interval (REI) = 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Beans (fresh & dry except succulent shelled)	Rust (<i>Uromyces appendiculatus</i>)	4 to 6 fl. oz. per acre (0.11 to 0.17 lb ai per acre)	Apply Axill Solutions Teb 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).
<p>Restrictions:</p> <p>Maximum Single Application Rate: 6 fl. oz. (0.17 lb ai) per acre Axill Solutions Teb 3.6SC.</p> <p>Maximum Annual Application Rate (Beans, fresh): 24 fl. oz. (0.68 lb ai) per acre per year Axill Solutions Teb 3.6SC. DO NOT make more than 6 applications per year when applying at the low rate.</p> <p>Maximum Annual Application Rate (Beans, dry): 12 fl. oz. (0.34 lb ai) per acre per year Axill Solutions Teb 3.6SC. DO NOT make more than 3 applications per year when applying at the low rate.</p> <p>Minimum Retreatment Interval (RTI): 14 days</p> <p>Pre-Harvest Interval: Beans, fresh: Axill Solutions Teb 3.6SC may be applied up to 7 days before harvest. Bean, dry: Axill Solutions Teb 3.6SC may be applied up to 14 days before harvest.</p> <p>Restricted-entry interval (REI) = 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Corn[*] (sweet corn, field corn, field corn grown for seed, and popcorn) [*Not approved for use on corn in New York State.]	Rust (<i>Puccinia</i> spp.) Northern leaf blight (<i>Helminthosporium turcicum</i>) Southern leaf blight (<i>Helminthosporium maydis</i>) Northern leaf spot (<i>Helminthosporium carbonum</i>) Gray leaf spot (<i>Cercospora zeae-maydis</i>)	4 to 6 fl. oz. per acre (0.11 to 0.17 lb ai per acre)	Apply Axill Solutions Teb 3.6SC in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).
<p>Restrictions:</p> <p>Maximum Single Application Rate: 6 fl. oz. (0.17 lb ai) per acre Axill Solutions Teb 3.6SC.</p> <p>Maximum Annual Application Rate: 24 fl. oz. (0.68 lb ai) per acre per year Axill Solutions Teb 3.6SC. DO NOT make more than 6 applications per year when applying at the low rate.</p> <p>Minimum Retreatment Interval (RTI): 7 days</p> <p>Pre-Harvest Interval (PHI): Sweet corn: Axill Solutions Teb 3.6SC may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: Axill Solutions Teb 3.6SC may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.</p> <p>Restricted-entry interval (REI) for sweet corn = 19 days</p> <p>Restricted-entry interval (REI) for all corn except sweet corn = 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Cotton	Southwestern cotton rust (<i>Puccinia cacabata</i>)	6 to 8 fl. oz. per acre (0.17 to 0.22 lb ai per acre)	<p>Apply Axill Solutions Teb 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control.</p> <p>For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p>
<p>Restrictions:</p> <p>Maximum Single Application Rate: 8 fl. oz. (0.22 lb ai) per acre Axill Solutions Teb 3.6SC.</p> <p>Maximum Annual Application Rate: 24 fl. oz. (0.68 lb ai) per acre per year Axill Solutions Teb 3.6SC. DO NOT make more than 4 applications per year when applying at the low rate.</p> <p>Minimum Retreatment Interval (RTI): 7 days</p> <p>Pre-Harvest Interval (PHI): Axill Solutions Teb 3.6SC may be applied up to 30 days before harvest.</p> <p>Restricted-entry interval (REI) = 12 hours</p>			

SPECIMEN

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd, (hyotan, cucuzza, hechima and Chinese okra) <i>Momordica</i> spp. (balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (butternut squash, calabaza,	Powdery mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)	4 to 6 fl. oz. per acre (0.11 to 0.17 lb ai per acre)	Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).
	Gummy stem blight – suppression (<i>Didymella bryonae</i>) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre (0.22 lb ai per acre)	

hubbard squash, acorn squash and spaghetti squash) Watermelon			
<p>Restrictions:</p> <p>Maximum Single Application Rate: 8 fl. oz. (0.22 lb ai) per acre Axill Solutions Teb 3.6SC.</p> <p>Maximum Annual Application Rate: 24 fl. oz. (0.68 lb ai) per acre per year Axill Solutions Teb 3.6SC.</p> <p>DO NOT make more than 6 applications per year when applying at the low rate.</p> <p>Minimum Retreatment Interval (RTI): 10 days</p> <p>Pre-Harvest Interval (PHI): Axill Solutions Teb 3.6SC may be applied up to 7 days before harvest.</p> <p>Restricted-entry interval (REI) = 12 hours</p>			

SPECIMEN

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Dry bulb onion Garlic Great-headed (elephant) garlic Welsh onion Shallot	White rot (<i>Sclerotium cepivorum</i>)	White rot: 20.5 fl oz (0.58 lb ai) per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.	<p>White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application must be made at the rate of 20.5 fl. oz. Axill Solutions Teb 3.6SC (0.58 lb ai) per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre.</p> <p>Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz. Axill Solutions Teb 3.6SC per acre (0.11 to 0.17 lb ai per acre) per application. Repeat at an interval of 10 to 14 days. Apply Axill Solutions Teb 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p>For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p>
	Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>) Purple blotch (<i>Alternaria porri</i>)	4 to 6 fl. oz. per acre (0.11 to 0.17 lb ai per acre)	
<p>Restrictions:</p> <p>Maximum Single Application Rate: 6 fl. oz. (0.17 lb ai) per acre Axill Solutions Teb 3.6SC unless applying for White rot in which case the maximum single application rate is 20.5 fl. oz. (0.58 lb ai) per acre.</p> <p>Maximum Annual Application Rate: 32.5 fl. oz. (0.91 lb ai) per acre per year Axill Solutions Teb 3.6SC if an in-furrow treatment is made. If not applied as an in-furrow treatment than the maximum annual application rate is 12 fl. oz. (0.34 lb ai) per acre per year Axill Solutions Teb 3.6SC as a foliar spray.</p> <p>DO NOT make more than 1 in-furrow application per year. If an in-furrow application is made, up to 2 additional foliar applications per year may be made. If no in-furrow application is made, DO NOT make more than 3 foliar applications per year when applying at the low rate.</p> <p>Minimum Retreatment Interval (RTI): 10 days</p> <p>Pre-Harvest Interval (PHI): DO NOT apply within 7 days of harvest.</p> <p>Restricted-entry interval (REI)= 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Fruiting Vegetables Group[*] Eggplant Groundcherry Pepino Pepper Tomatillo Tomato [*Not registered for this use in California.]	Early blight (<i>Alternaria solani</i>)	8 fl. oz. per acre (0.22 lb ai per acre)	Apply Axill Solutions Teb 3.6SC as a foliar spray using an interval of 7 days. For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).
<p>Restrictions:</p> <p>Maximum Single Application Rate: 8 fl. oz. (0.22 lb ai) per acre Axill Solutions Teb 3.6SC.</p> <p>Maximum Annual Application Rate: 48 fl. oz. (1.35 lb ai) per acre per year Axill Solutions Teb 3.6SC. DO NOT make more than 6 applications per year.</p> <p>Minimum Retreatment Interval (RTI): 7 days</p> <p>Pre-Harvest Interval (PHI): DO NOT apply within 7 days of harvest.</p> <p>Restricted-entry interval (REI) = 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Grasses Grown for Seed	Rusts (<i>Puccinia</i> spp.)	4 to 8 fl. oz. per acre (0.11 to 0.22 lb ai per acre)	<p>Apply the specified rate of Axill Solutions Teb 3.6SC as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure, use 6 to 8 fl. oz./A (0.17 to 0.22 lb ai/A) and shorter spray intervals. Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.</p> <p>For optimum benefit, tank mix the lowest specified rate of a spray surfactant with Axill Solutions Teb 3.6SC.</p>
	Powdery mildew	4 to 8 fl. oz. per acre (0.11 to 0.22 lb ai per acre)	<p>Apply specified rate of Axill Solutions Teb 3.6SC when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./A (0.17 to 0.22 lb ai/A) and shorter spray intervals. Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.</p> <p>For optimum benefit, tank mix the lowest specified rate of a spray surfactant with Axill Solutions Teb 3.6SC.</p>
<p>Restrictions:</p> <p>Chaff, screenings and straw from treated areas may be used for feed purposes; however, DO NOT forage, cut green chop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.</p> <p>Maximum Single Application Rate: 8 fl. oz. (0.22 lb ai) per acre Axill Solutions Teb 3.6SC.</p> <p>Maximum Annual Application Rate: 16 fl. oz. (0.45 lb ai) per acre per year Axill Solutions Teb 3.6SC. DO NOT make more than 4 applications per year when applying at the low rate.</p> <p>Minimum Retreatment Interval (RTI): 14 days</p> <p>Pre-Harvest Interval (PHI): Axill Solutions Teb 3.6SC may be applied up to 4 days before harvest.</p> <p>Restricted-entry interval (REI) = 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (<i>Sclerotium cepivorum</i>) suppression only Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>) Purple blotch (<i>Alternaria porri</i>)	4 to 6 fl. oz. per acre (0.11 to 0.17 lb ai per acre)	For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Axill Solutions Teb 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development. For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).
<p>Restrictions:</p> <p>Maximum Single Application Rate: 6 fl. oz. (0.17 lb ai) per acre Axill Solutions Teb 3.6SC.</p> <p>Maximum Annual Application Rate: 24 fl. oz. (0.68 lb ai) per acre per year Axill Solutions Teb 3.6SC.</p> <p>DO NOT make more than 6 applications per year when applying at the low rate.</p> <p>Minimum Retreatment Interval (RTI): 10 days</p> <p>Pre-Harvest Interval (PHI): DO NOT apply within 7 days of harvest.</p> <p>Restricted-entry interval (REI) = 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Hops	Powdery mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i>)	4 to 8 fl. oz. per acre (0.11 to 0.22 lb ai per acre)	Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Increase the spray volume and the application rate as vine growth increases during the season. For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

Maximum Single Application Rate: 8 fl. oz. (0.22 lb ai) per acre Axill Solutions Teb 3.6SC.
Maximum Annual Application Rate: 32 fl. oz. (0.90 lb ai) per acre per year Axill Solutions Teb 3.6SC.
DO NOT make more than 8 applications per year when applying at the low rate.
Minimum Retreatment Interval (RTI): 10 days
Pre-Harvest Interval (PHI): Axill Solutions Teb 3.6SC may be applied up to 14 days before harvest.
Restricted-entry interval (REI) = 12 hours

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (<i>Cercospora brassicicola</i>) Powdery mildew (<i>Erysiphe cruciferarum</i>) Alternaria leaf spot (<i>Alternaria brassicicola</i>)	3 to 4 fl. oz. per acre (0.08 to 0.11 lb ai per acre)	For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions: Application to turnip greens is limited to East of the Rockies.

Maximum Single Application Rate: 4 fl. oz. (0.11 lb ai) per acre Axill Solutions Teb 3.6SC.
Maximum Annual Application Rate: 16 fl. oz. (0.45 lb ai) per acre per year Axill Solutions Teb 3.6SC.
DO NOT make more than 5 applications per year when applying at the low rate.
Minimum Retreatment Interval (RTI): 10 days
Pre-Harvest Interval (PHI): **DO NOT** apply within 7 days of harvest.
Restricted-entry interval (REI) = 12 hours

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Garden beet roots and tops (leaves)	Cercospora leaf spot (<i>Cercospora beticola</i>)	3 to 7.2 fl. oz. per acre (0.08 to 0.20 lb ai per acre)	Make applications on a 14 day interval. For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3)
<p>Restrictions:</p> <p>Maximum Single Application Rate: 7.2 fl. oz. (0.20 lb ai) per acre Axill Solutions Teb 3.6SC. Maximum Annual Application Rate: 28.8 fl. oz. (0.81 lb ai) per acre per year Axill Solutions Teb 3.6SC. DO NOT make more than 9 applications per year when applying at the low rate. Minimum Retreatment Interval (RTI): 14 days Pre-Harvest Interval (PHI): DO NOT apply within 7 days of harvest. Restricted-entry interval (REI) = 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Lychee	Anthrachnose (<i>Colletotrichum gloeosporioides</i>)	4 to 6 fl. oz. per acre (0.11 to 0.17 lb ai per acre)	Begin first application of Axill Solutions Teb 3.6SC as panicle emerges. Spray up to 6 fl. oz. (0.17 lb ai) per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. For optimum disease control, tank mix the lowest labeled rate of a non-ionic spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).
<p>Restrictions:</p> <p>Maximum Single Application Rate: 6 fl. oz. (0.17 lb ai) per acre Axill Solutions Teb 3.6SC. Maximum Annual Application Rate: 48 fl. oz. (1.35 lb ai) per acre per year Axill Solutions Teb 3.6SC. DO NOT make more than 8 applications per year.</p>			

Minimum Retreatment Interval (RTI): 10 days
Pre-Harvest Interval (PHI): Axill Solutions Teb 3.6SC can be applied up to and including the day of harvest (PHI = 0 days).
Restricted-entry interval (REI) = 2 days

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Okra	Cercospora leaf spot (<i>Cercospora</i> spp.)	4 to 6 fl. oz. per acre (0.11 to 0.17 lb ai per acre)	Apply specific dosage of Axill Solutions Teb 3.6SC in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:
Maximum Single Application Rate: 6 fl. oz. (0.17 lb ai) per acre Axill Solutions Teb 3.6SC.
Maximum Annual Application Rate: 24 fl. oz. (0.68 lb ai) per acre per year Axill Solutions Teb 3.6SC.
DO NOT make more than 6 applications per year when applying at the low rate.
Minimum Retreatment Interval (RTI): 14 days
Pre-Harvest Interval (PHI): **DO NOT** apply within 3 days of harvest.
Restricted-entry interval (REI) = 12 hours

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Peanut	<p>SOILBORNE: Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only)</p> <p>FOLIAR: Early leaf spot Late leaf spot Leaf rust Web blotch (<i>Phoma</i>) Pepper spot (<i>Leptoshaerulina</i>)</p>	7.2 fl. oz. per acre (0.20 lb ai per acre)	<p>FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventative spray schedule. Applications of chlorothalonil must be made prior to and following applications of Axill Solutions Teb 3.6SC to discourage development of resistant strains of fungi. For optimum control of foliar diseases including leaf rust, web blotch, and pepper spot, tank mix the lowest label specified rate of a spray surfactant with Axill Solutions Teb 3.6SC.</p> <p>LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Axill Solutions Teb 3.6SC in the first advisory spray in July and continue Axill Solutions Teb 3.6SC applications at 14-day intervals. Applications after August 15 must be tank mixed with chlorothalonil for resistance management purposes. For optimum control of the specified soilborne diseases, four consecutive applications of Axill Solutions Teb 3.6SC must be made at 14-day intervals. Axill Solutions Teb 3.6SC is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Axill Solutions Teb 3.6SC as a leaf spot resistance management strategy. A spray surfactant is not necessary when Axill Solutions Teb 3.6SC is tank mixed with chlorothalonil. Mixing or alternating Axill Solutions Teb 3.6SC with other DMI fungicides may lead to resistance.</p> <p>Axill Solutions Teb 3.6SC must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i>. Drought conditions will decrease the effectiveness of Axill Solutions Teb 3.6SC against the root and pod rots.</p> <p>Use Axill Solutions Teb 3.6SC in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, for example, proper crop rotation practices.</p>
<p>Restrictions: DO NOT feed hay or threshings or allow livestock to graze in treated areas.</p> <p>Maximum Single Application Rate: 7.2 fl. oz. (0.20 lb ai) per acre Axill Solutions Teb 3.6SC.</p>			

Maximum Annual Application Rate: 28.8 fl. oz. (0.81 lb ai) per acre per year Axill Solutions Teb 3.6SC.

DO NOT make more than 4 applications per year.

Minimum Retreatment Interval (RTI): 14 days

Pre-Harvest Interval (PHI): Axill Solutions Teb 3.6SC may be applied up to 14 days before harvest.

Restricted-entry interval (REI) = 12 hours

Timing of Axill Solutions Teb 3.6SC Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	Axill Solutions Teb 3.6SC Application No.	Chlorothalonil Application No.
7 Applications	3,4,5, and 6	1, 2 and 7

SPECIMEN

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Pecan	Brown leaf spot <i>(Sirosporium diffusum)</i> Downy spot <i>(Mycosphaerella caryigena)</i> Liver spot <i>(Gnomonia caryae)</i> Scab <i>(Cladosporium caryigenum)</i> Vein spot <i>(Gnomonia nerviseda)</i> Zonate leaf spot <i>(Grovesinia pyramidalis)</i>	4 to 8 fl. oz. per acre (0.11 to 0.22 lb ai per acre)	<p>Apply Axill Solutions Teb 3.6SC in a preventative spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Axill Solutions Teb 3.6SC can be applied at 4 fl. oz. (0.11 lb ai) per acre in a tank-mix with the specified rate of Super-Tin® in cover sprays. Follow label directions for the use of Super-Tin. DO NOT add surfactant to the spray solution when tank-mixing Axill Solutions Teb 3.6SC with Super-Tin. Apply Axill Solutions Teb 3.6SC in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. (0.20 to 0.22 lb ai) per acre of Axill Solutions Teb 3.6SC to full-size mature trees, and 4 to 6 fl. oz. (0.11 to 0.17 lb ai) per acre of Axill Solutions Teb 3.6SC to smaller trees. Apply the highest specified rate to varieties that are highly susceptible to the indicated diseases, or when severe diseases conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases.</p> <p>For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p>
<p>Restrictions:</p> <p>DO NOT apply after shucks begin to split.</p> <p>DO NOT cut cover crops in treated areas for feed or allow livestock to graze treated areas.</p> <p>Maximum Single Application Rate: 8 fl. oz. (0.22 lb ai) per acre Axill Solutions Teb 3.6SC.</p> <p>Maximum Annual Application Rate: 32 fl. oz. (0.90 lb ai) per acre per year Axill Solutions Teb 3.6SC.</p> <p>DO NOT make more than 8 applications per year when applying at the low rate.</p> <p>Minimum Retreatment Interval (RTI): 10 days</p> <p>Restricted-entry interval (REI) = 12 hours</p>			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Soybean	Rust (<i>Phakopsora pachyrhizi</i>) Powdery Mildew (<i>Microsphaera diffusa</i>)	3 to 4 fl. oz. per acre (0.08 to 0.11 lb ai per acre)	Apply Axill Solutions Teb 3.6SC as a broadcast foliar spray as a preventative spray or at first visible symptoms of diseases. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use the highest specified rates and shorter spray intervals when disease pressure is severe. The lowest labeled rate of spray surfactant must be tank-mixed with Axill Solutions Teb 3.6SC. Apply Axill Solutions Teb 3.6SC in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.
Restrictions: Maximum Single Application Rate: 4 fl. oz. (0.11 lb ai) per acre Axill Solutions Teb 3.6SC. Maximum Annual Application Rate: 12 fl. oz. (0.34 lb ai) per acre per year Axill Solutions Teb 3.6SC. DO NOT make more than 4 applications per year when applying at the low rate. Minimum Retreatment Interval (RTI): 10 days Pre-Harvest Interval (PHI): DO NOT apply within 21 days of harvest. Restricted-entry interval (REI) = 12 hours			

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Sunflower	Rust (<i>Puccinia helianthi</i>)	4 to 6 fl. oz. per acre (0.11 to 0.17 lb ai per acre)	Apply specific dosage of Axill Solutions Teb 3.6SC at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply highest specified rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. For optimum disease control, tank mix the lowest labeled rate of a spray surfactant should be tank-mixed with Axill Solutions Teb 3.6SC. Contact your state Extension Service representative for a list of approved surfactants. Axill Solutions Teb 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).
Restrictions:			

Maximum Single Application Rate: 6 fl. oz. (0.17 lb ai) per acre Axill Solutions Teb 3.6SC.
Maximum Annual Application Rate: 16 fl. oz. (0.45 lb ai) per acre per year Axill Solutions Teb 3.6SC.
DO NOT make more than 4 applications per year when applying at the low rate.
Minimum Retreatment Interval (RTI): 14 days
Pre-Harvest Interval (PHI): 50 days
Restricted-entry interval (REI) = 12 hours

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Turnip (Application is limited to East of the Rockies)	Cercospora leaf spot (<i>Cercospora brassicicola</i>)	4 to 7.2 fl. oz. per acre (0.11 to 0.20 lb ai per acre)	Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:
Maximum Single Application Rate: 7.2 fl. oz. (0.20 lb ai) per acre Axill Solutions Teb 3.6SC.
Maximum Annual Application Rate: 28.8 fl. oz. (0.81 lb ai) per acre per year Axill Solutions Teb 3.6SC.
DO NOT make more than 7 applications per year when applying at the low rate.
Minimum Retreatment Interval (RTI): 12 days
Pre-Harvest Interval (PHI): 7 days
Restricted-entry interval (REI) = 12 hours

CROP	DISEASE	RATE OF AXILL SOLUTIONS TEB 3.6SC	USE DIRECTIONS
Wheat	Rusts leaf, stem, and stripe (<i>Puccinia</i> spp.) Head blight or scab (<i>Fusarium</i> spp.) - Suppression	4 fl. oz. per acre (0.11 lb ai per acre)	<p>Observe wheat fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Apply Tebuconazole 3.6C in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air.</p> <p>For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Axill Solutions Teb 3.6SC. Axill Solutions Teb 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Axill Solutions Teb 3.6SC will be resistant to weathering. Axill Solutions Teb 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Application Timing Directions: Rusts: Apply Axill Solutions Teb 3.6SC at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Axill Solutions Teb 3.6SC for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p>
<p>Restrictions: Straw may be fed or used for bedding. DO NOT allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Axill Solutions Teb 3.6SC.</p> <p>Maximum Single Application Rate: 4 fl. oz. (0.11 lb ai) per acre Axill Solutions Teb 3.6SC DO NOT make more than one application per year.</p> <p>Pre-Harvest Interval (PHI): DO NOT apply within 30 days of harvest.</p> <p>Restricted-entry interval (REI) = 12 hours</p>			

<p>SEED TREATMENT: Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn) For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.</p>
<p>The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:</p> <ul style="list-style-type: none"> • This seed has been treated with Axill Solutions Teb 3.6SC fungicide containing tebuconazole. • DO NOT use treated seed for food, feed or oil purposes. <p>The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with tebuconazole:</p> <ul style="list-style-type: none"> • Store treated seed away from food and feedstuffs. • DO NOT allow children, pets or livestock to have access to treated seed. • Wear long pants, long-sleeved shirt, shoes plus socks, and protective gloves when handling treated seed. • Treated seed exposed on soil surfaces may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting (for example, in row ends). • Dispose of all excess treated seed by burying seed away from bodies of water.

- **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

USE RESTRICTIONS: Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 153.155(c).

DISEASE	RATE Fl Oz/CWT	DIRECTIONS FOR USE
Soilborne and Seedborne Fusarium	0.071 (0.002 lb ai)	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed must be sound and well cured prior to treatment. Dilute product with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates advised for the crop to be treated with Axill Solutions Teb 3.6SC. The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut (<i>Sphacelotheca reilana</i>)	0.27 – 0.54 (0.008 – 0.015 lb ai)	

SPECIMEN

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

DO NOT apply by ground or air within 100 feet of aquatic areas listed above.

DO NOT cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

DISEASE CONTROL IN GOLF COURSE TURF, FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

Chemigation: **DO NOT** apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES

- **DO NOT** apply within 100 feet of the aquatic areas listed above.
- **DO NOT** cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for further information.
- **DO NOT** apply when wind velocity exceeds 15 mph.
- **DO NOT** make ground applications during temperature inversions.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Spray Volume: For best results Axill Solutions Teb 3.6SC may be applied in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are advised depending upon the equipment, plant species and plant growth stage at the time of application. For the most effective results, check equipment calibration regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1 ½ teaspoon for each pint of liquid product of specified label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution must not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label.

Mixing: Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that Axill Solutions Teb 3.6SC is completely dispersed in the mixture.

Mixing Instructions: **DO NOT** allow the ratio of water:product in field to exceed 1:1. Fill the spray tank (or chemigation nurse tank) half full with water before adding any product. **DO NOT** add product to the spray tank before adding water, this may result in a hazardous chemical reaction.

Mixing Procedure:

1. Water. Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2. Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3. Inductor. If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA Bags. Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water Dispersible Products. Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6. Water-soluble products.
7. Emulsifiable concentrates (including oil concentrates when applicable).
8. Water soluble additives (including AMS or UAN when applicable)
9. Remaining quantity of water.

DISEASE CONTROL IN GOLF COURSE TURF

Turf Use Restrictions

- For use on golf course turf only.
- **DO NOT** use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
- Not for residential use.
- Not for use on turf being grown for sale or commercial use as sod.
- **DO NOT** use clippings for animal feed.
- **Maximum Single Application Rate:** 3.6 fl. oz. of Axill Solutions Teb 3.6SC (0.10 lb ai) per 1,000 sq ft per year.
- **Minimum Retreatment Interval (RTI):** 28 days
- **DO NOT** apply more than 6 applications per year.
- [In New York State **DO NOT** apply more than 3 applications of 0.6 fl. oz. of Axill Solutions Teb 3.6SC not to exceed 1.8 fl. oz. of product per 1,000 sq. ft. per year.]
- [In the State of New York, **DO NOT** exceed 1.8 fl. oz. of this product per 1,000 sq. ft. (2.2 lbs. a.i./A) per year.]

Product Information

For use on all golf turf applications of cool season and warm season grasses (including Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Axill Solutions Teb 3.6SC is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Bermudagrass can be sensitive to Axill Solutions Teb 3.6SC under certain conditions. **DO NOT** apply consecutive applications during or just after dormancy break. Avoid application when temperatures are expected to exceed 85 degrees F.

Axill Solutions Teb 3.6SC can be used for the prevention and control of the diseases mentioned in the table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of Axill Solutions Teb 3.6SC in sufficient water for thorough coverage. A volume of 66-132 gallons per acre (1.5-3.0 gallons per 1,000 sq. ft.) is advised. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, Axill Solutions Teb 3.6SC must be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE of Axill Solutions Teb 3.6SC (Fl. oz/1000 Sq Ft)	USE DIRECTIONS AND RESTRICTIONS
Dollar Spot (<i>Sclerotinia homeocarpa</i>) Copper Spot (<i>Gloeocercospora sorghi</i>) Powdery Mildew (<i>Erysiphe graminis</i>) Corticium Red Tread (<i>Laetisaria fuciformis</i>) Rusts (<i>Puccinia</i> spp.)	0.6-1.1 (0.017-0.031 lb ai)	For prevention, begin applications when conditions are favorable for disease development. Restrictions: DO NOT make two consecutive applications of Axill Solutions Teb 3.6SC. Alternate with another fungicide with a different mode of action. A second application may be made after 28 days.

DISEASE	RATE of Axill Solutions Teb 3.6SC (Fl. oz/1000 Sq Ft)	USE DIRECTIONS AND RESTRICTIONS
Brown Patch/Rhizoctonia Blight, Large Patch (<i>Rhizoctonia solani</i>) Brown Ring Patch (<i>R. circinata</i>)	0.6-1.1 (0.017-0.031 lb ai)	For prevention, begin applications when conditions are favorable for disease development. Restrictions: DO NOT make two consecutive applications of Axill Solutions Teb 3.6SC. Alternate with a different mode of action. A second application may be made after 28 days.
Anthracnose – Basal and Foliar (<i>Colletotrichum cereal</i>) Red Thread (<i>Laetisaria fuciformis</i>) Pink Patch (<i>Limonomyces roseipellis</i>)	0.6-1.1 (0.017-0.031 lb ai)	
Bermuda Grass decline (<i>Gaeumannomyces graminis var. graminis</i>)	0.6-1.1 (0.017-0.031 lb ai)	Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent of the depth of the root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Restrictions: Apply subsequent applications at 28 day intervals.
Take All Patch (<i>Gaeumannomyces graminis</i>)	0.6-1.1 (0.017-0.031 lb ai)	For prevention, apply in the fall when soil temperature reaches 55-65°F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Restrictions: Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.6-1.1 (0.017-0.031 lb ai)	Apply when conditions are favorable for disease development at 28 day intervals.
Stipe Smut (<i>Ustilago striiformis</i>)	0.6-1.1 (0.017-0.031 lb ai)	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot (<i>Leptosphaeria korrae</i> , <i>L. narmari</i> , <i>Ophiosphaerella herpotricha</i> , <i>Gaeumannomyces graminis</i>) Necrotic Ring Spot (<i>Leptosphaeria korrae</i>)	0.6-1.1 (0.017-0.031 lb ai)	For prevention, apply in fall when soil temperature reaches 65°F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (<i>Fusarium roseum</i>)	0.6-1.1 (0.017-0.031 lb ai)	Apply first application in mid-June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 28 day intervals.

DISEASE	RATE of Axill Solutions Teb 3.6SC (Fl. oz/1000 Sq Ft)	USE DIRECTIONS AND RESTRICTIONS
Summer Patch (<i>Magnaporthe poae</i>)	0.6-1.1 (0.017-0.031 lb ai)	Apply beginning in the spring. See local university guidance for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone. Restrictions: DO NOT make two consecutive applications of Axill Solutions Teb 3.6SC. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28 day intervals.
Zoysia Patch, Large Patch of zoysia (<i>Rhizoctonia solani</i>)	0.6-1.1 (0.017-0.031 lb ai)	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/Typhula Blight (<i>Typhula incarnate</i>) Pink Snow Mold/Microdochium Patch (<i>Microdochium nivalis</i>)	0.6-1.1 (0.017-0.031 lb ai)	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. It is advised the Axill Solutions Teb 3.6SC be tank-mixed with other registered snow mold products for best season long results. Restrictions: DO NOT apply over snow cover, or when turf is dormant.
Restrictions: Apply the specified amount of Axill Solutions Teb 3.6SC in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. DO NOT use clippings for animal feed. DO NOT exceed 3.6 fl. oz. of Axill Solutions Teb 3.6SC (0.10 lb ai) per 1000 sq. ft. per year. DO NOT exceed 6 applications per year.		

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS, AND ORNAMENTALS IN COMMERCIAL AND RESIDENTIAL LANDSCAPES

Ornamental Use Restrictions:

- For use on ornamental plants only; not for use on woodlands or forest management.
- **DO NOT** apply more than 10 fl. oz. (0.28 lb ai) per acre in a single application.
- **DO NOT** apply more than 0.31 gallon (40 fl. oz.) of Axill Solutions Teb 3.6SC (equal to 1.13 lbs. of tebuconazole) per acre per year.
- **DO NOT** make more than 4 applications per year.
- **Minimum Retreatment Interval (RTI):** 14 days; except for applications to prevent Petal Blight where applications may be made 2-3 times per week into the flowers as they open and develop color.
- **Restricted-entry interval (REI):** 12 hours
- **DO NOT** apply to bearing fruit trees or vegetables.

Axill Solutions Teb 3.6SC can be used in a preventative and curative disease control program for the listed plant types and diseases in the table below. Optimum disease management is obtained when Axill Solutions Teb 3.6SC is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin application when disease first appears and repeat at 14-21 day intervals during the growing season. Use

the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply Axill Solutions Teb 3.6SC at rates of 4-10 fl. oz. (0.11-0.28 lb ai) per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The “Directions for Use” of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. **DO NOT** use this product on African Violets, Begonias, Boston Fern, and Geraniums.

Ornamentals Disease Control

PLANTS	DISEASE	USE DIRECTIONS	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or Tank mixing with barrier protectant fungicides is advised for resistance management.	
Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthrachnose Leaf Spot Powdery Mildew Rust Scab		
Azaleas, Camellas, Rhododendrons, and Other Landscape (Ornamental) Shrubs	Anthrachnose Black Spot Leaf Spot Petal Blight	Petal Blight – Apply 2-3 times per week into the flowers as they open and develop color.	
Ground Covers and Vines	Powdery Mildew Rust Southern Blight		
HOW MUCH TO USE FOR SMALL PLANTINGS: ADD 1 TEASPOON TO 2.5 GALLONS OF WATER.			

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the spray tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust the nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store above 28°F or agitate before use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: [Note to reviewer: One of the optional Container Handling options will appear on the final printed label.]

[Nonrefillable plastic containers less than or equal to 5 gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

[Nonrefillable plastic containers greater than 5 gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

[Refillable Containers greater than 55 gallons: Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.]

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Axill Solutions, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Axill Solutions, LLC and Seller harmless for any claims relating to such factors.

Axill Solutions, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Axill Solutions, LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, Axill Solutions, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Axill Solutions, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF AXILL SOLUTIONS, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF AXILL SOLUTIONS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Axill Solutions, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Axill Solutions, LLC.

SPECIMEN